

Systems for Sparse Representations: Fourier Analysis, Wavelets & Shearlets

Felix Voigtlaender

DEDALE Tutorial Day
November 16, 2016




1 Wavelets

2 Curvelets & Shearlets


1 Wavelets

2 Curvelets & Shearlets

Definition: Let $\phi, \psi \in L^2(\mathbb{R})$ be a scaling function and a wavelet (i.e., $\int_{\mathbb{R}} |\hat{\psi}(\omega)|^2 \frac{d\omega}{|\omega|} < \infty$). Then the **inhomogeneous discrete wavelet system** generated by ϕ, ψ is

$$\left(\phi(\bullet - m) \right)_{m \in \mathbb{Z}} \cup \left(2^{j/2} \cdot \psi(2^j \bullet - m) \right)_{j \in \mathbb{N}_0, m \in \mathbb{Z}}.$$


Definition: Let $\phi, \psi \in L^2(\mathbb{R})$ be a scaling function and a wavelet (i.e., $\int_{\mathbb{R}} |\hat{\psi}(\omega)|^2 \frac{d\omega}{|\omega|} < \infty$). Then the **inhomogeneous discrete wavelet system** generated by ϕ, ψ is


$$\left(\phi(\bullet - m) \right)_{m \in \mathbb{Z}} \cup \left(2^{j/2} \cdot \psi(2^j \bullet - m) \right)_{j \in \mathbb{N}_0, m \in \mathbb{Z}}.$$


Definition: Let ϕ, ψ as above. A **2D inhomogeneous discrete wavelet system** is defined by

$$\left(\phi^{(1)}(\bullet - m) \right)_{m \in \mathbb{Z}^2} \cup \left(2^j \cdot \psi^{(\ell)}(2^j \bullet - m) \right)_{j \in \mathbb{N}_0, m \in \mathbb{Z}^2, \ell \in \{1, 2, 3\}}$$

where

$$\phi^{(1)} := \phi \otimes \phi, \quad \psi^{(1)} := \phi \otimes \psi, \quad \psi^{(2)} := \psi \otimes \phi, \quad \psi^{(3)} := \psi \otimes \psi.$$




Approximation properties of Wavelets

For good **compression** of a signal f , it would be desirable that f can be well approximated using only a few wavelets:


Approximation properties of Wavelets

For good **compression** of a signal f , it would be desirable that f can be well approximated using only a few wavelets:



Approximation properties of Wavelets

For good **compression** of a signal f , it would be desirable that f can be well approximated using only a few wavelets:




Theorem: Discrete wavelets provide **optimal approximation rates** for those $f \in L^2(\mathbb{R}^2)$ which are C^2 apart from **point singularities**:

$$\|f - f_N\|_{L^2} \lesssim N^{-1/2} \quad (N \rightarrow \infty).$$

Living on the point





Living on the point edge



Natural images are governed by curved singularities, not point singularities!

Living on the point edge



Natural images are governed by curved singularities, not point singularities!

Definition (Donoho; 2001)

With $Q := (0,1)^2$, the set of **cartoon-like functions** is defined as

$$\mathcal{E}^2(\mathbb{R}^2) = \{f_0 + f_1 \cdot \mathbf{1}_B \mid \partial B \subset Q \text{ closed } C^2 \text{ curve and } f_0, f_1 \in C_c^2(Q)\}.$$

Approximation of cartoon-like functions

Theorem: For $f \in \mathcal{E}^2(\mathbb{R}^2)$, the best N -term approximation f_N

- using **Fourier basis**: $\|f - f_N\|_{L^2} \lesssim N^{-1/4}$,
- using **Wavelets**: $\|f - f_N\|_{L^2} \lesssim N^{-1/2}$.

Approximation of cartoon-like functions

Theorem: For $f \in \mathcal{E}^2(\mathbb{R}^2)$, the best N -term approximation f_N

- using **Fourier basis**: $\|f - f_N\|_{L^2} \lesssim N^{-1/4}$,
 - using **Wavelets**: $\|f - f_N\|_{L^2} \lesssim N^{-1/2}$.
-

Theorem (Donoho; 2001): Let $(\psi_n)_{n \in \mathbb{N}}$ be **any** countable family in $L^2(\mathbb{R}^2)$. If

$$\|f - f_N\|_{L^2} \lesssim N^{-\theta} \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2),$$

then $\theta \leq 1$. Here, we assume **polynomial depth search** for forming f_N .

Approximation of cartoon-like functions

Theorem: For $f \in \mathcal{E}^2(\mathbb{R}^2)$, the best N -term approximation f_N

- using **Fourier basis**: $\|f - f_N\|_{L^2} \lesssim N^{-1/4}$,
- using **Wavelets**: $\|f - f_N\|_{L^2} \lesssim N^{-1/2}$.

Theorem (Donoho; 2001): Let $(\psi_n)_{n \in \mathbb{N}}$ be **any** countable family in $L^2(\mathbb{R}^2)$. If

$$\|f - f_N\|_{L^2} \lesssim N^{-\theta} \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2),$$

then $\theta \leq 1$. Here, we assume **polynomial depth search** for forming f_N .

Hence, Wavelets **cannot** approximate curvilinear singularities optimally.

Approximation of cartoon-like functions

Theorem: For $f \in \mathcal{E}^2(\mathbb{R}^2)$, the best N -term approximation f_N

- using **Fourier basis**: $\|f - f_N\|_{L^2} \lesssim N^{-1/4}$,
- using **Wavelets**: $\|f - f_N\|_{L^2} \lesssim N^{-1/2}$.

Theorem (Donoho; 2001): Let $(\psi_n)_{n \in \mathbb{N}}$ be **any** countable family in $L^2(\mathbb{R}^2)$. If

$$\|f - f_N\|_{L^2} \lesssim N^{-\theta} \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2),$$

then $\theta \leq 1$. Here, we assume **polynomial depth search** for forming f_N .

Hence, Wavelets **cannot** approximate curvilinear singularities optimally.

Intuitive explanation: This is caused by the **scalar** dilations:



1 Wavelets

2 Curvelets & Shearlets



Towards a directional system

Recall: **Isotropic** dilations are suboptimal for approximating curves:




Towards a directional system

Recall: Isotropic dilations are suboptimal for approximating curves:



Idea for shearlets: Use

- Parabolic scaling
- Different orientations via shearing.




Recall: Isotropic dilations are suboptimal for approximating curves:



Idea for shearlets: Use

- Parabolic scaling
- Different orientations via shearing.



Advantages of shearing:

- The shearing matrices $\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$ leave the digital grid \mathbb{Z}^2 invariant.
- Uniform theory for the analog and digital situation.

A first definition of shearlets

Definition (Kutyniok, Labate; 2006): For $\psi \in L^2(\mathbb{R}^2)$, the associated discrete shearlet system is


$$\left(2^{\frac{3}{4}j} \cdot \psi(S_k A_{2^j} \bullet - m) \right)_{j,k \in \mathbb{Z}, m \in \mathbb{Z}^2}.$$

A first definition of shearlets


Definition (Kutyniok, Labate; 2006): For $\psi \in L^2(\mathbb{R}^2)$, the associated discrete shearlet system is

$$\left(2^{\frac{3}{4}j} \cdot \psi(S_k A_{2^j} \bullet - m) \right)_{j,k \in \mathbb{Z}, m \in \mathbb{Z}^2}.$$


Induced frequency tiling:



The induced frequency tiling




The induced frequency tiling




Very different treatment of x -direction and y -direction!

The induced frequency tiling




Very different treatment of x -direction and y -direction!

Solution: Use cone-adapted shearlets:





The induced frequency tiling




Very different treatment of x -direction and y -direction!

Solution: Use cone-adapted shearlets:





The induced frequency tiling



Very different treatment of x -direction and y -direction!

Solution: Use cone-adapted shearlets:



Cone-adapted Discrete Shearlet Systems

Definition (Kutyniok, Labate; 2006):

The **cone-adapted discrete shearlet system** $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ with **sampling density** $c > 0$ generated by $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ is the union of


$$\{\phi(\bullet - c \cdot m) \mid m \in \mathbb{Z}^2\},$$

$$\{\psi_{j,k,m,h} := \psi(S_k A_{2^j} \bullet - c \cdot m) \mid j \in \mathbb{N}_0, |k| \leq \lceil 2^{j/2} \rceil, m \in \mathbb{Z}^2\},$$

$$\{\psi_{j,k,m,v} := \tilde{\psi}(S_k^T \tilde{A}_{2^j} \bullet - c \cdot m) \mid j \in \mathbb{N}_0, |k| \leq \lceil 2^{j/2} \rceil, m \in \mathbb{Z}^2\},$$



where

$$S_k = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}, \quad A_{2^j} = \begin{pmatrix} 2^j & 0 \\ 0 & 2^{j/2} \end{pmatrix}, \quad \tilde{A}_{2^j} = \begin{pmatrix} 2^{j/2} & 0 \\ 0 & 2^j \end{pmatrix}.$$



Curvelets induce a similar frequency tiling than shearlets, but using rotations instead of shearings.

Curvelets induce a similar frequency tiling than shearlets, but using rotations instead of shearings.



Theorem (Kittipoom, Kutyniok, Lim; 2012):

Let $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ be compactly supported and assume that

- $\widehat{\psi}, \widehat{\tilde{\psi}}$ satisfy certain decay conditions,
- we have

$$|\widehat{\phi}(\xi)|^2 + \sum_{j,k} |\widehat{\psi}_{j,k}(\xi)|^2 + |\widehat{\tilde{\psi}}_{j,k}(\xi)|^2 \geq C > 0 \quad \text{a.e.} \quad (\dagger)$$

Then there is some $c > 0$ such that $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ is a frame for $L^2(\mathbb{R}^2)$.

Compactly supported shearlets


Theorem (Kittipoom, Kutyniok, Lim; 2012):

Let $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ be compactly supported and assume that

- $\widehat{\psi}, \widehat{\tilde{\psi}}$ satisfy certain decay conditions,
- we have

$$|\widehat{\phi}(\xi)|^2 + \sum_{j,k} |\widehat{\psi}_{j,k}(\xi)|^2 + |\widehat{\tilde{\psi}}_{j,k}(\xi)|^2 \geq C > 0 \quad \text{a.e.} \quad (\dagger)$$

Then there is some $c > 0$ such that $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ is a frame for $L^2(\mathbb{R}^2)$.



Theorem (Kittipoom, Kutyniok, Lim; 2012):

Let $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ be compactly supported and assume that

- $\widehat{\psi}, \widehat{\tilde{\psi}}$ satisfy certain decay conditions,
- we have

$$|\widehat{\phi}(\xi)|^2 + \sum_{j,k} |\widehat{\psi}_{j,k}(\xi)|^2 + |\widehat{\tilde{\psi}}_{j,k}(\xi)|^2 \geq C > 0 \quad \text{a.e.} \quad (\dagger)$$

Then there is some $c > 0$ such that $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ is a frame for $L^2(\mathbb{R}^2)$.

Remarks

- For (\dagger) , it suffices to have $\tilde{\psi}((\xi_1, \xi_2)) = \psi((\xi_2, \xi_1))$ as well as

$$\begin{aligned} |\widehat{\phi}(\xi)| &\gtrsim 1 && \text{for } \xi \in [-1, 1]^2, \\ |\widehat{\psi}(\xi)| &\gtrsim 1 && \text{for } \xi_1 \in [1/3, 3] \text{ and } |\xi_2| \leq |\xi_1|, \end{aligned}$$

- There are special examples with frame bounds $B/A \approx 4$.

Theorem (Kutyniok, Lim; 2011):

Let $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ be compactly supported and such that $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ is a frame for $L^2(\mathbb{R}^2)$. Further, assume that $\hat{\psi}, \hat{\tilde{\psi}}$ satisfy certain decay conditions. Then $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ provides an **optimally sparse approximation** of all $f \in \mathcal{E}^2(\mathbb{R}^2)$, i.e.,

$$\|f - f_N\|_{L^2} \lesssim N^{-1} \cdot (\log N)^{3/2} \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2).$$

Theorem (Kutyniok, Lim; 2011):

Let $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ be compactly supported and such that $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ is a frame for $L^2(\mathbb{R}^2)$. Further, assume that $\hat{\psi}, \hat{\tilde{\psi}}$ satisfy certain decay conditions. Then $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ provides an **optimally sparse approximation** of all $f \in \mathcal{E}^2(\mathbb{R}^2)$, i.e.,

$$\|f - f_N\|_{L^2} \lesssim N^{-1} \cdot (\log N)^{3/2} \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2).$$

Here, f_N is a linear combination of (at most) N elements of the **dual frame** of $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$.

Theorem (Kutyniok, Lim; 2011):

Let $\phi, \psi, \tilde{\psi} \in L^2(\mathbb{R}^2)$ be compactly supported and such that $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ is a frame for $L^2(\mathbb{R}^2)$. Further, assume that $\hat{\psi}, \hat{\tilde{\psi}}$ satisfy certain decay conditions. Then $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$ provides an **optimally sparse approximation** of all $f \in \mathcal{E}^2(\mathbb{R}^2)$, i.e.,

$$\|f - f_N\|_{L^2} \lesssim N^{-1} \cdot (\log N)^{3/2} \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2).$$

Here, f_N is a linear combination of (at most) N elements of the **dual frame** of $\mathcal{SH}(\phi, \psi, \tilde{\psi}; c)$.

Remark

The proof shows

$$\sum_{n>N} |\theta(f)|_n^2 \lesssim N^{-2} \cdot (\log N)^3 \quad \forall N \in \mathbb{N} \quad \forall f \in \mathcal{E}^2(\mathbb{R}^2),$$

where $|\theta(f)|_n$ denotes the n -th largest shearlet coefficient. Hence, the shearlet coefficients are in ℓ^p for all $p > \frac{2}{3}$.

Thank you!

Thank you!

Questions, comments, counterexamples?

